GT121

GL Sciences Inc.


Enrichment Analysis of Volatile Components of Edible Oils — Comparisons of Sampling Methods using MonoTrap and HandyTD

Using a trapping media MonoTrap RGC18 TD and a thermal desorption device HandyTD TD265, we conducted a simplified screen-analysis of volatile components arising from pseudo-degraded oils (heated for 8 h in 180 °C oven).

This time, the samples were heated at 60 °C and sampled by headspace gas sampling method. Thermostatic Chamber Static method which means leave the sample into oven and heating. Bath Shaking method means heating while shaking in a water bath at a constant temperature. After a comparison of both methods it found the bath shaking method was able to detect volatile components with higher sensitivity.

Furthermore, used the bath shaking method to compare the differences of volatile compounds between edible oil just after opening and the degraded oil. Some typical oil odor compounds such as 2, 4-Heptadienal, 2, and 4-Decadienal were detected with good sensitivity from degraded edible oil.

Sample Preperation Procedure



GC/MS Conditions

System: Thermal Desorption-GC/MS

(HandyTD TD265)

Column : InertCap Pure-WAX

 $0.25 \text{ mm I.D. X } 60 \text{ m, df} = 0.5 \text{ }\mu\text{m}$

Col. Cat. No. : 1010-68164

Col. Temp. : 40 °C(5 min) - 5 °C/min - 250 °C Carrier Gas : He, 1 mL/min (constant flow)

GC Inlet : 250 °C Split 10:1 **Detection** : MS Scan (*m/z* 30-350)


HandyTD Conditions

Desorb Temp. : Room temperature- 45 °C/sec-

250 °C (5 min)


Pre Desorb Press.: 140 kPa

Comparison of Sampling Methods (degraded oil)



Qualitative of ingredients from library searches can be found on the following pages

Comparison of Oil Just After Opening and Degraded oil (water bath shaking method)



Library search results.

- 2. Pentanal
- 3. 3-Penten-2-one
- 4. 6-Ethyl-2-methyldecane
- 5. 1-Penten-3-one
- 6. 2-Butenal
- 7. 6-Methyl-1-heptanol
- 8. 4-Hexen-3-one
- 9. Hexanal
- 10. 2-Pentenal
- 11. 1-Penten-3-ol
- 12. 2-Heptanone

- 13. Heptanal
- 14. 3-Hexen-2-one
- 15. 2-Hexenal
- 16. 2-Pentylfuran
- 17. 1-Pentanol
- 18. 2-Octanone
- 19. Octanal
- 20. Ethylhexanol
- 21. 2-Heptenal
- 22. Nonanal

GL Sciences B.V.

5652 AS Eindhoven

Phone: +31 (0)40 254 95 31

Email: info@glsciences.eu

www.glsciences.eu

The Netherlands

De Sleutel 9

- 23. Acetic acid
- 24. 2-Octenall

- 25. 1-Octen-3-ol
- 26. Heptanol
- 27. 2,4-Heptadienal
- 28. 2,4-Heptadiena
- 29. Propanoic acid
- 30. Benzaldehyde
- 31. 2-Nonenal
- 32. 1-Octanol
- 33. Butanoic acid
- 34. Butyrolactone
- 35. 2-Decenal 36. Vinyl caproate

- 37. Pentanoic acid 38. 2-Undecenal
- 39. 2,4-Decadienal
- 40. 2,4-Decadienal
- 41. Hexanoic acid
- 42. Heptanoic acid

GL Sciences disclaims any and all responsibility for any injury or damage which may be caused by this data directly or indirectly. We reserve the right to amend this information or data at any time and without any prior announcement.

GL Sciences, Inc. Japan

22-1 Nishishinjuku 6-Chome Shinjuku-ku, Tokyo,

Phone: +81-3-5323-6620 +81-3-5323-6621

163-1130, Japan

Email: world@gls.co.jp

Web: www.glsciences.com

International Distributors

Visit our Website at:

https://www.glsciences.com/company/distributor.html

GL Sciences (ShangHai) Ltd.

Tower B, Room 2003, Far East International Plaza, NO,317 Xianxia Road, Changning District.

Shanghai, China P.C. 200032

Phone: +86 (0)21-6278-2272 Email: contact@glsciences.com.cn

www.glsciences.com.cn

Email: info@glsciencesinc.com www.glsciencesinc.com

310-265-4425

GL Sciences, Inc. USA

Torrance, CA 90503

Fax:

Phone: 310-265-4424

4733 Torrance Blvd. Suite 255

